

PROLONGED PREMEDICATION IN ELECTIVE SURGERY: PROSPECTS AND LIMITATIONS

S.Sh. Joniev, T. Yuldashev, I. Shodmonov

¹Samarkand State Medical University, Samarkand, Uzbekistan

Abstract. The article presents the results of preoperative preparation, anesthesia and surgical treatment of patients operated on for non-toxic nodular goiter intensive care, acute coronary syndrome. A new approach to preoperative preparation for thyroid surgery is described. The effectiveness in the preoperative period of using the modified method of preoperative preparation using sibazon and droperidol and anesthesia with the use of ketamine and the advantages of this method compared with other methods of general anesthesia are shown.

Key words: intensive care, acute coronary syndrome modified preoperative preparation, anesthesia, thyroid gland, nontoxic nodular goiter

Acute coronary syndrome., there is a steady increase in the number of thyroid diseases worldwide. A significant number of population living in the territory of Uzbekistan have obvious or hidden functional disorders of the thyroid gland [6]. Diffuse non - toxic goiter is the most common pathology, which takes up to 60% of all cases of thyroid disease. In this pathology, the functions of the central nervous system and endocrine system, blood circulation and respiration, liver and kidneys, immunity and metabolism are impaired [5]. Often, this type of thyroid disease is the leading one in the group of endocrine diseases, the main method of treatment for which is surgery. It should be considered that during operations on the thyroid gland, it is important to use the optimal method of anesthesia, which would prevent the manifestations of pathological reactions associated with the nature of the main and concomitant diseases. The high risk of intra - and postoperative complications associated with the anatomical

features of the surgical intervention area justifies the relevance of the problem of optimizing the anesthetic allowance for thyroid surgery [2].

Analyzing of currently used methods of general anesthesia for thyroid diseases, we can assume that not all of them do fully prevent the negative effects and reactions that occur in the body to surgical stress and have many other serious drawbacks. These include: the use of narcotic analgesics and anesthetics, postoperative respiratory depression and rapid cessation of analgesia in the early postoperative period, a number of adverse hemodynamic changes at traumatic stages of surgery [1].

When choosing an anesthetic to maintain anesthesia for thyroid disease, the characteristics of the psychological and somatic status of patients, the nature of the influence of the disease on the circulatory system and the functional state of parenchymal organs, and the presence of concomitant diseases are guided [3].

Surgical treatment of thyroid pathology, including goiter under General anesthesia using neuroleptanalgesia (NLA) in the most traumatic moments of the operation is often accompanied by dangerous circulatory disorders in the form of tachycardia, arterial hypertension, and heart rhythm disorders [3]. If the patient also has concomitant diseases (CVS diseases and diabetes mellitus), then the complication during anesthesia becomes critical. In modern practical anesthesiology, much attention is paid to the blockage of pathological impulsion, which occurs under the influence of surgical trauma of the afferent and central nervous system during the medical preparation of the patient in the preoperative period.

Anesthesiology does not yet know ideal and universal solutions to the problem of protecting the patient from surgical aggression. The most reasonable approach is a multi-modal approach that implies a multi-level, multi-purpose antinociception, in which the maximum effect (due to synergy or summation of action) is combined with a minimum of side effects [6].

The appearance of modern drugs for anesthesia and improvement of surgical treatment results at the current stage of development of endocrine surgery is seen in the further improvement of preoperative preparation and intraoperative anesthesia.

Objective: To choose the effective method of preoperative preparation and types of anesthesia on acute coronary syndrome and thyroid gland surgery.

Material and methods:

The research was conducted at Samarkand regional branch of the Republican specialized scientific practical medical center of Cardiology. 72 patients operated on non-toxic goiter were under observation. In accordance with the purpose and objectives of this study, patients were divided into two groups depending on the type of preoperative preparation and anesthesia. Among the examined patients there were 8 men (11.1%) and 64 women (88.9%) aged from 32 to 68 years. By age, the patients were distributed as follows: from 32-45 years – 13 people (18.05 %), 46-60 years – 49 people (74.7 %), over 60 years – 10 people (7.2%). The length of anamnesis for goiter was on average 3.3 ± 2 years. Objective status according to the classification of the American society of anesthesiologists (ASA) II - 39 (54, 1%), III - 28 (38.9%), IV - 5 (6.9%). Patients with nodular (multi-nodular) euthyroid colloid goiter were operated on.

The following operations were performed: strumectomy (14 cases), hemistrumectomy (24 cases), hemistrumectomy with isthmus removal (7 cases), extremely subtotal-subfacial strumectomy (11 cases). The average duration of the operation is 50 ± 13 minutes. The first group (control group - $n=34$) – patients who underwent traditional preoperative therapy and standard anesthesia. Group II (study group - $n=38$) - patients whose preoperative preparation was performed using a modified method with the use of sibazone and droperidol. In group 1, premedication was performed on the operating table: fentanyl 0.002 mg/kg, sibazone 5 mg, atropine 0.005-0.008 mg/kg. Initial narcosis — thiopental Na 4 -7 mg/kg. Intubation was

performed on the background of mioplegii ditilinom (100mg). To maintain anesthesia, propofol 2 — 4 mg/kg/h, fentanyl 5 — 8 mcg kg/h, and droperidol 0.05 — 0.1 mg/kg were used. Patients in group 2 were given sibazone 0.2-0.5 mg/kg at 20:00 I/M for 3 days before surgery. In addition to the standard premedication, sibazone at a dose of 0.3-0.5 mg/kg and droperidol 0.05-0.1 mg/kg were administered 30 to 40 minutes before surgery. Introductory anesthesia — thiopental Na 4 — 7 mg/kg. Intubation on the background of myoplegia with ditillin(100 mg). To maintain anesthesia, propofol 2 — 4 mg/ kg/h, fentanyl 3 — 5 mcg/ kg/h, droperidol 0.05 — 0.1 mg/kg, ketamine 0.5 mg/kg were used. To objectively evaluate the effectiveness of preoperative preparation and the adequacy of anesthesia, hemodynamic parameters were studied: systolic blood pressure (SBP, mmHg), diastolic blood pressure (DBP, mmHg), heart rate (HR, beats/min) were determined in dynamics by the " ARGUS TM-7 "monitor of the company "SCHILLER". Average dynamic blood pressure (ABP, mmHg) $SBP = DBP + 1/3 (SBP-DBP)$ (B. Folkov, E. Neil, 1976). The concentration of glucose, lactate, SpO₂, and hormonal parameters (cortisol, free T₃, and TSH) were studied using the STAR-FAX immunoassay analyzer (USA). The level of sedation was determined on the Ramsay scale (M. A. Ramsay, 1974) 40 minutes after premedication. The study of hemodynamic parameters was performed five times: at admission, 2 days to operation, 1 day to operation, in the intraoperative period, on the 1st day after surgery.

RESEARCH RESULT

Our studies showed that the initial parameters of central hemodynamics in patients in both groups did not significantly differ from each other (tables 1., 2.). conducting a step-by-step monitoring of changes in central hemodynamics, we found that patients in the control group already at the preoperative stage, before induction into anesthesia, there was a significant increase in blood pressure, SBP, DBP, ABP and heart rate ($p < 0.05$) compared to the initial parameters. So, after premedication, patients in the control group showed a significant increase in blood pressure by 4.8% ($p < 0.001$), SBP by 6.9% ($p < 0.001$), ABP by 5.5% ($p < 0.01$), heart rate by 4.4% ($p < 0.05$) relative to the

initial values (table 1.). the number of heart contractions, average blood pressure during the three days before surgery was steadily increased and despite the traditional antihypertensive therapy, there was no downward trend. It is also noteworthy that despite the traditional premedication, the number of heartbeats was increased compared to the previous days.

The above data indicate that patients in the control group have significant changes in blood pressure and heart rate, which are a consequence of the impact on the patient's body of stress and other adverse factors acting on the patient's body in the perioperative period. These disorders are not completely blocked by premedication, anesthesia, or infusion therapy and are amplified under the influence of surgery. Analysis of central hemodynamic parameters in patients of the study group showed that in the preoperative period, at the first five stages of the study (3 days, 2 days, 1 day before surgery, premedication), against the background of the use of sibazone and droperidol, there was a systematic decrease in blood pressure, SBP, DBP, heart rate compared to the initial indicators, but within the physiological norm. 2 days before the operation, there was a significant decrease in SBP by 4.2% ($p < 0.005$), DBP by 4.3% ($p < 0.01$), ABP by 4.2% ($p < 0.01$), and heart rate by 3.9% ($p < 0.05$) compared to the first stage. After premedication, the SBP is lower than the initial figures by 3.4% ($p < 0.01$), DBP by 5.3% ($p < 0.001$), ABP by 4.5% ($p < 0.001$), heart rate by 4.6% ($p < 0.05$). These changes in central hemodynamic parameters are positive and are due to the stabilization of the neurovegetative system against the background of the use of sibazone and droperidol, since admission to the hospital itself is already a stressful situation for most patients.

By analyzing the level of preoperative sedation, it was found that in 80% of patients in the control group, the effect of premedication was unsatisfactory, it was expressed in emotional tension, anxiety, and fear of surgery. In the study group, the level of preoperative sedation was adequate in 100 % of cases.

At the traumatic stage of the operation, there was a significant increase in the average blood pressure values in group 1 by 19.2 % ($p < 0.05$), in group 2-by 12 % (p

< 0.05). Heart rate in the most traumatic stages of surgery increased by 15.6 % ($p < 0.05$) in group 1 and by 16 % ($p < 0.05$) in group 2. These changes indicated a hyperdynamic reaction of the cardiovascular system, activation of the neuro-vegetative system. There were no significant differences between the two groups at this stage of the study ($p > 0.05$). SBP returned to normal in group 2 after the operation, and in group 1 only by the first day after the operation. In the postoperative period, the heart rate remained stable.

In group 1, the glucose level increased during the traumatic stage of the operation, reaching a maximum by the end of the operation (6.98 mmol/l; $p < 0.05$), and returned to normal only on the first day. In group 2, the blood glucose concentration was normalized by the first day after surgery.

The level of TSH and T3 in both groups remained within the reference values at all stages of the study, no significant differences in these indicators were found in the comparison groups ($p > 0.05$). In all groups, SpO₂ remained at the normal level of 97-99% during anesthesia and in the early postoperative period.

Conclusions:

1. Patients operated on for thyroid diseases, in the intraoperative period, undesirable hemodynamic, vegetative and neuroendocrine reactions of the body occur, which negatively affect the course of the perioperative period and the anesthetic allowance.
2. The use of a modified method of preoperative preparation with the use of sibazone and droperidol in patients operated on for thyroid diseases helps to reduce emotional stress, providing an adequate level of preoperative sedation, allows you to optimize the anesthetic effects, minimize the negative effects and doses of anesthetics.

3. The use of general anesthesia with fentanyl and droperidol does not fully block nociceptive impulses, which indicates insufficient protection of the patient from surgical aggression, characterized by instability of hemodynamics, preservation of endocrine and metabolic changes. The addition of ketamine to General anesthesia and the use of a modified method of preoperative preparation can reduce the dose of opioids, stabilize hemodynamics and ensure safety of the perioperative period.

Literature:

1. 1. Aralov U.A. Joniev, S., Rakhimov, Improving the effectiveness of preoperative preparation methods in thyroid surgery [Text] Aralov, S. Sh. Rakhimov // Problemy biologii i meditsiny. — 2015. — No. 1. — S. 11-14.
2. Joniev S.Sh., Rakhimov A.U., Comparison of methods of preoperative preparation in thyroid gland / Joniev S.Sh., Rakhimov A.U., [Текст] // VI International conference "Sharing the results of research towards closer global convergence of scientists". — Ontario, Canada: 1807-150 Charlton av. East Hamilton, 2015. — С. 38-43.
3. 3. Joniev, S. Sh., Babajanov, A. S. Assessment of pre-operative preparation and effectiveness of anesthesia in thyroid pathology operations / S. Sh. Joniev, A. S. Babajanov [Text] // Proceedings of the Samarkand regional scientific practical conference "XXI century - the century of the intellectual generation". — Samarkand, Uzbekistan: SamDChTI publishing house, 2016. — S. 190-193.
4. Joniev S.Sh. Features of preoperative preparation and anesthesia in thyroid pathology [Текст] / Joniev S.Sh. // British Medical Journal . — 2022. — № Volume-2, No 4. — С. 212-215.
5. Joniev S.Sh. Anesthesiological allowance for operations for diffuse goiter [Текст] / Joniev S.Sh. // Journal of integrated education and research. — 2022. — № Volume 1, Issue 5. — С. 19-26.

6. Joniev S.Sh. Improving the results of anesthesia in thyroid pathology [Текст] / Joniev S.Sh. // Uzbek medical journal. — 2022. — № Volume 3, Issue 3. — С. 23-28.
7. Joniev S. Sh., Pardaev Sh. K. Use of general multicomponent anesthesia in thyroid operations [Text] / S. Sh. Joniev, Sh. K. Pardaev // A new day in medicine. — 2021. — No. 6 (38/1). — С. 443-479. Joniev S.Sh. Determining the level of preoperative preparation and conducting anesthesia in patients with thyroid pathologies [Текст] / Joniev S.Sh. // Danish Scientific Journal (DSJ). — 2022. — № 59. — С. 19-23.
8. Joniev S.Sh. Improvement of the results of anesthesia in thyroid pathology [Текст] / Joniev S.Sh. // Annali d'Italia. — 2022. — № Vol 1. 30. — С. 78-82.
9. Joniev S.Sh., Yakubov I., Pormonov Kh., Daminov I., Ensuring adequate premedication in patients with thyroid pathology [Текст] / Joniev S.Sh., Yakubov I., Pormonov Kh., Daminov I., // Norwegian Journal of development of the International Science. — 2022. — № 86. — С. 17-20.
10. Joniev S.Sh., Tukhsanbaev S., Kurbanov K., Abdaliev D., Predicting the effectiveness of anesthesia in thyroid pathology [Текст] / Joniev S.Sh., Tukhsanbaev S., Kurbanov K., Abdaliev D., // Polish journal of science. — 2022. — № 51. — С. 30-34.
11. Joniev S.Sh., Ganiev A., Ibrokhimov Z., Melikboboev A., Analysis of the effectiveness of anesthesia methods in thyroid pathology [Текст] / Joniev S.Sh., Ganiev A., Ibrokhimov Z., Melikboboev A., // Znanstvena misel journal Slovenia. — 2022. — № 67. — С. 32-35.
12. Joniev S.Sh., Pardaev Sh.K., Muhammadieva U., Analysis of methods of general anesthesia in thyroid surgery based on hemodynamic parameters [Текст] / Joniev S.Sh., Pardaev Sh.K., Muhammadieva U., // Turkish Journal of Physiotherapy and Rehabilitation. — 2021. — № 32(3). — С. 7136-7140.

13. Joniev S.Sh., Pardaev Sh.K., Akramov B.R., Hushvakov U.O., Monitoring And Evaluation Of Hemodynamic Parameters During Anesthesia In Endocrine Surgery [Текст] / Joniev S.Sh., Pardaev Sh.K., Akramov B.R., Hushvakov U.O., // The American Journal of Medical Sciences and Pharmaceutical Research. — 2020. — № 2(12). — С. 40-46.
14. Joniev S.Sh., Hemodynamic aspects of multicomponent general anesthesia and endocrine surgery [Text] / Joniev S.Sh., // MEDICUS International Medical Scientific Journal. — 2020. — No. No. 5 (35). — S. 8-13.
15. Joniev, S. Sh., Pardaev, Sh. K., Goyibov, S. S., Akramov, B. R. Use of a modified method of preoperative preparation and anesthesia and surgery of the thyroid gland / S. Sh. Joniev, Sh. K. Pardaev, S. S. Goyibov, B. R. Akramov [Text] // Collection of scientific articles XIV International correspondence scientific specialized conference. — Boston: PROBLEMS OF SCIENCE, 2019. — S. 175-186.
16. Joniev S. Sh., Khushvaktov U. O. Rational hemodynamic monitoring of anesthesiological procedures and endocrine surgery // Actual diagnostic questions and treatment of novel coronavirus infection. - 2020. - S. 6-11.
17. Joniev, S. Sh., Muminov, A. A., Khushvaktov, U. O. Peri operative period and endocrine surgery: puti reshenia anesthesiologicheskikh problem [Text] / S. Sh. Joniev, A. A. Muminov, U. O. Khushvaktov // Eurazisky Soyuz Uchenykh. — 2020. — No. 5-6 (74). — S. 38-41.
18. Joniev S. Sh., Rakhimov A. U. Varianty preoperatsionnoy podgotovki pri operatsiyax na shchitovidnoy jjele [Text] / Joniev S. Sh., Rakhimov A. U. // Nauka i mir. — 2014. — No. 11-2. — S. 138-141.
19. Joniev S. Sh., Influence of complex preoperative preparation and biochemical indicators on blood of patients with nodular zobos [Text] / Joniev S. Sh., // Doctor's newsletter. — 2013. — No. 1(3). — S. 71-73.

20. Joniev S.Sh, Rakhimov, A. U., Babajanov, A. S. Znachenie biochemical pokazatelei pri preoperatsionnoy pregotovki bolnykh uzlovym zobom [Text] / A. U. Rakhimov, A. S. Babajanov // Science and world. — 2013. — No. 9(136). — S. 136-138.
21. Joniev S. Sh., Babajanov A. S., Khushnaev S., Sultanova S. Uluchshenie metodov preoperatsionnoy pregotovki i anestheziy v perioperatsionnom periode zbolevani schitovidnoy izlezny [Text] / Joniev S. Sh., Babajanov A. S., Khushnaev S., Sultanova S. // European research. — 2018. — No. 5. — S. 135-138.
22. Joniev S. Sh., Babajanov A. S., Rakhimov, A. U. Analysis of the effectiveness of variants of preoperative preparation and anesthesia and treatment of pain in the thyroid gland [Text] / Joniev S. Sh., Babajanov A. S., A. U. Rakhimov // Problemy biology and medicine. — 2017. — No. 1. — S. 93-95.
23. Joniev S.Sh., Determining the effectiveness of preoperative preparation of patients with nodular zobos and biochemical indicators. [Text] / S. Sh. Joniev // Problemy biology and medicine. — 2014. — No. 1 (77). — S. 14-18.
24. Joniev, S. Sh. Modified method of preoperative preparation and anesthesia in operative treatment of pathology of schitovidnoy signs / S. Sh. Joniev [Text] // Materialy mejregionalnogo nauchno-prakticheskogo foruma s mejdunarodnym uchastiem "Zdorove semi — budushchee Rossii". — Izhevsk: OOO "MedEkspoForum", 2019. — S. 37-40.
25. Joniev, S. Sh. Pre-operative preparation and anesthesia during operation with no shield / S. Sh. Joniev [Text] // Sbornik tsikla lektsii "Anesteziya u pasitinov s soputstvuyushchimi zabolevaniyami i pri razlichnykh tipakh khirurgicheskikh vmeshatelstv". — Samarkand: SamGMI, 2015. — P. 72-81.

26. Rakhimov, A. U., Joniev, S. Sh. Assessment of the effectiveness of general anesthesia methods in thyroid pathology operations [Text] / A. U. Rakhimov, S. Sh. Joniev // Problemy biology and medicine. — 2022. — No. 4,1(138). — S. 118-125. IL Sharipov, JT Yusupov, BK Xolbekov Personalization and preventative premedication: used drugs value and efficiency //Web of Scientist: International Scientific Research Journal 3 (02), 740-748
27. Matlubov Mansur Muratovich, Yusupov Jasur Tolibovich, Mallayev Surat Sadullayevich, Khamrayev Khamza Hamidullayevich Optimization of anesthesiological assistance in women with arterial hypertension in hysterectomy // Достижения науки и образования. 2020. №5 (59).
28. Muratovich, Matlubov Mansur, et al. "Hemodynamic indicators in pregnant women with obesity of various degrees of expression." *European Journal of Molecular and Clinical Medicine*, vol. 8, no. 2, 15 Jan. 2021